

Influence of $SU(2) \otimes U(1)$ singlet scalars on Higgs boson signal at LHC

N.V.Krasnikov *

CERN, Geneva 23, Switzerland

September,1997

Abstract

We investigate a model of $SU(2) \otimes U(1)$ singlet scalars coupled to Standard Model. We point out that for the case of the maximal mixing between singlet scalars and standard Higgs boson the Higgs boson production cross section is smaller than the standard Higgs boson production by factor $\frac{1}{N+1}$, where N is the number of singlet scalars. For the case of big mixing Higgs boson could be nonobservable at LHC. However, there is also a possibility for the observation of scalar singlets at LHC. As a quasirealistic example we discuss supersymmetric $SU(2) \otimes U(1)$ electroweak model with an additional singlet chiral superfield.

*On leave of absence from INR, Moscow 117312, Russia

One of the main tasks of existing and future collider experiments (LEP2, LHC, NLC,...) is the search for the Higgs boson. Its detection would be the confirmation of the spontaneous symmetry breaking mechanism and an experimental "proof" of the renormalizability of the electroweak interactions. For the two most popular models of electroweak physics, the Standard Model(SM) and its minimal supersymmetric generalization(MSSM) LEP2 and LHC will be able to discover the Higgs boson with a mass from 77 GeV(current standard Higgs boson mass limit [1]) up to 10^3 GeV(upper limit comes from requirement of tree level unitarity [2]). In the SM and MSSM the Higgs boson for $m_h \leq 400$ GeV (this bound comes from an analysis of high precision LEP1 data [3]) is rather narrow that leads to sharply peaked resonance. The signal to background ratio in the resonance region is mostly determined by the experimental energy resolution.

In this paper we investigate the influence of $SU(2) \otimes U(1)$ singlet scalar fields on the detection of the Higgs boson at LHC. Namely, we investigate the influence of the mixing between $SU(2) \otimes U(1)$ scalar singlets and the Higgs boson. We find that in an extreme case of maximal mixing between $SU(2) \otimes U(1)$ scalars and the Higgs boson the Higgs boson production cross section drops by factor $\frac{1}{N+1}$ (N is the number of scalar singlets) that could lead to nonobservability of the Higgs boson at LEP or LHC. For instance, for LHC for $N \geq 2$ it is possible to make Higgs signal nonobservable for an integrated luminosity $L = 10^5 pb^{-1}$. However, there is also possibility for the observation of the scalar singlets at LHC. As a quasirealistic example we discuss a supersymmetric electroweak model with singlet chiral superfield. In the limit when the number of singlet scalar fields $N \rightarrow \infty$ we find that the Higgs boson mass is continuously distributed. A model of the Higgs boson with continuously distributed mass has been proposed in ref. [4]. It should be noted that the models with additional light $SU(2) \otimes U(1)$ singlet scalar boson have been discussed in the literature [5]. In such models Higgs boson invisible decay modes into light singlet scalars result in a modification of branching ratios that leads to slightly different strategy for the Higgs boson search. In recent paper of T.Binoth and J.J. van der Bij [6] a model of light scalar singlets strongly coupled to the standard Higgs boson has been discussed. In such model Higgs boson decays mainly into light singlet scalar bosons and its decay

width is not small that considerably affects the signal to background ratio S/B making Higgs signal diluted. We consider slightly different situation when big mixing of the Higgs boson with scalar singlets results in the appearance of many "Higgs like" scalar bosons in the spectrum and the production cross section of "Higgs like" bosons is smaller than the standard Higgs boson cross section production that leads to the Higgs signal dilution. In considered model the Higgs bosons branching ratios are not modified compared to the standard Higgs boson case.

Let us start with the simplest modification of the SM model, namely, we add to the SM an additional real scalar singlet $\phi(x)$. Take the effective potential in the form

$$V = \lambda(H^+H - \frac{v^2}{2})^2 + k(H^+H - \frac{v^2}{2})\phi + \frac{M^2\phi^2}{2}. \quad (1)$$

After the spontaneous symmetry breaking $\langle H \rangle = \frac{v}{\sqrt{2}}$, $\langle \phi \rangle = 0$ we find that in the unitary gauge $H(x) = (0, \frac{v+h(x)}{\sqrt{2}})$ the mass matrix for the Higgs field $h(x)$ and singlet fields $\phi(x)$ takes the form

$$\hat{M} = m_h^2 h^2(x) + M^2 \phi^2(x) + 2\delta m^2 h(x)\phi(x), \quad (2)$$

where $m_h^2 = 2\lambda v^2$ and $\delta m^2 = \frac{kv}{2}$. The eigenstates of the mass matrix (2) are the fields

$$h_1(x) = h(x) \cos(\delta) - \phi(x) \sin(\delta), \quad (3)$$

$$\phi_1(x) = \phi(x) \cos(\delta) + h(x) \sin(\delta) \quad (4)$$

with eigenvalues

$$M_{h_1, \phi_1} = \frac{m_h^2 + M^2}{2} \pm \sqrt{\frac{(m_h^2 - M^2)^2}{4} + (\delta m^2)^2} \quad (5)$$

and with mixing angle

$$\tan(2\delta) = \frac{2\delta m^2}{M^2 - m_h^2}. \quad (6)$$

Original Higgs field $h(x)$ is expressed in terms of the $h_1(x)$, $\phi_1(x)$ fields as

$$h(x) = h_1(x) \cos(\delta) + \phi_1(x) \sin(\delta) \quad (7)$$

The main production mechanisms of the Higgs boson at LEP and LHC are the Higgs bremsstrahlung of a real(LEP2) or virtual(LEP1) Z-boson and gluon fusion correspondingly. As a consequence of the formula (7) we find that the production cross sections of the h_1 and ϕ_1 bosons are the same as for the standard Higgs boson production with the corresponding mass except multiplicative factors $\cos^2(\delta)$ (h_1 -boson) and $\sin^2(\delta)$ (ϕ_1 -boson). The branching ratios of the h_1 and ϕ_1 bosons into fermion-antifermion pairs, 2 gluons or 2 photons are the same as for the SM Higgs boson. As a result of the additional factors $\cos^2(\delta)$, $\sin^2(\delta)$ the signal significance $S = \frac{N_S}{\sqrt{N_B}}$ will be smaller than for the case of the SM Higgs boson provided the mass difference between the h_1 and ϕ_1 bosons is bigger than the detector mass resolution. To be concrete consider the Higgs boson production at LHC ¹. For the case $100 \text{ GeV} \leq m_h \leq 130 \text{ GeV}$ the most promising signature for the search for the Higgs boson is

$$pp \rightarrow (h \rightarrow \gamma\gamma) + \dots \quad (8)$$

For low luminosity stage with an integral luminosity $L = 3 \cdot 10^4 pb^{-1}$ CMS detector will be able to discover the standard Higgs boson with the signal significance [8] $6 \leq S \leq 9$. The ATLAS detector [9] will have the similar signal significance as the CMS. The mass resolution of the diphoton pair invariant mass at low luminosity stage is assumed to be $\delta m_{\gamma\gamma} \approx 0.5 \text{ GeV}$. For the case of maximal mixing $\delta = \frac{\pi}{4}$ and for $M_{h_1} - M_{\phi_1} \geq 1 \text{ GeV}$ as a result of nonzero mixing we shall have two resonances in the diphoton spectrum. For the case of maximal mixing the Higgs boson production cross section decreases by factor 2 and the signal significance of each of two Higgs like resonances will be between 3 and 4.5. According to standard convention Higgs boson is assumed to be discovered provided the signal significance is bigger than 5. So at low luminosity stage in the case of maximal mixing the Higgs boson escapes from being discovered. At high luminosity stage for an integral luminosity $L = 10^5 pb^{-1}$ the signal significance for CMS detector for $110 \text{ GeV} \leq m_h \leq 140 \text{ GeV}$ is estimated to be $11 \leq S \leq 13$. At high luminosity stage two Higgs like resonances will be detected at $S \geq 5$ significance level. Therefore there is

¹LEP1 data [7] lead to the bound $\sin^2(\delta) \leq 0.1$ for $m_{\phi_1} \leq 30 \text{ GeV}$

nonzero probability instead of the single Higgs boson to discover 2 Higgs bosons by the measurement of the diphoton spectrum.

For the Higgs boson mass interval $140 \text{ GeV} \leq m_h \leq 600 \text{ GeV}$ the most promising signature for the Higgs boson detection is through the decay

$$h \rightarrow ZZ^*(Z) \rightarrow 4\text{leptons}. \quad (9)$$

For the Higgs boson mass $200 \text{ GeV} \leq m_h \leq 400 \text{ GeV}$ the signal significance for an integrated luminosity $L = 10^5 \text{ pb}^{-1}$ is estimated to be $S = 16 - 18$ [8]. For such mass interval the maximal mixing will be detected at $S \geq 5$ significance level. Moreover for such mass interval it is possible to detect two Higgs like resonances at 5σ level provided the mixing angle $\sin^2(\delta) \geq 0.32$.

At LEP1 and LEP2 the effects of nonzero mixing are the same as at LHC provided the mass difference between the scalars h_1 and ϕ_1 is bigger than the detector energy resolution $\delta E \sim 4 \text{ GeV}$. Again instead of the single Higgs boson we shall have two Higgs like bosons with production cross sections for the maximal mixing two times smaller the standard one. Therefore the signal significance $S = \frac{N_S}{\sqrt{N_B}}$ is decreased by factor two compared to the standard case. An increase of an integral luminosity L by factor 4 just compensates the decrease of the Higgs boson production cross section.

Consider now the case of N additional $SU(2) \otimes U(1)$ singlet scalar fields. Take the effective potential in the form

$$V = \lambda(H^+H - \frac{v^2}{2})^2 + k_i(H^+H - \frac{v^2}{2})\phi_i + \frac{1}{2}m_{ij}^2\phi_i\phi_j, \quad (10)$$

where $i, j = 2, 3, \dots, N+1$. The equations for the determination of the nontrivial minimum of the effective potential (10) have solution $\langle H \rangle = \frac{v}{\sqrt{2}}$, $\langle \phi_i \rangle = 0$. In the unitare gauge $H = (0, \frac{v+h(x)}{\sqrt{2}})$ the mass matrix for the Higgs field $\phi_1(x) \equiv h(x)$ and singlet fields $\phi_i(x)$ ($i = 2, 3, \dots, N+1$) has the form

$$\hat{M}^2 = m_{ij}^2\phi_i(x)\phi_j(x), \quad (11)$$

where $i, j = 1, 2, \dots, N+1$, $m_{11}^2 = 2\lambda v^2$, $m_{1i}^2 = m_{i1}^2 = \frac{k_i v}{2}$. After the diagonalization the

mass matrix (11) takes the form

$$\hat{M}^2 = M_i^2 (\phi'_i)^2, \quad (12)$$

where $\phi'_i = O_{ij}\phi_j$, $\phi_i = O_{ji}\phi'_j$ and $O_{ij}O_{ik} = \delta_{jk}$. Parameters m_{ij}^2 are arbitrary so in general orthogonal matrix O_{ij} can have arbitrary form. Consider as an extreme example the case of maximal mixing, i.e.

$$O_{1,1} = O_{2,1} = \dots = O_{N+1,1} = \frac{1}{\sqrt{N+1}}. \quad (13)$$

For the mixing matrix (13) the original Higgs field $h(x) \equiv \phi_1(x)$ is a superposition of the of the fields $\phi'_i(x)$ with definite mass

$$h(x) = \frac{1}{\sqrt{N+1}} \sum_{i=1}^{N+1} \phi'_i \quad (14)$$

Again as in the case of $N = 1$ one can find that the branching ratios for the Higgs like bosons $\phi'_i(x)$ are the same as for the standard Higgs boson but their production cross section drops by factor $\frac{1}{N+1}$ compared to the standard Higgs boson case. For the Higgs boson mass range interval $100 \text{ GeV} \leq m_h \leq 130 \text{ GeV}$ and for an integrated luminosity $L = 10^5 \text{ pb}^{-1}$ the signal significance in an extreme case of maximal mixing is $2.7 \leq S \leq 4.1$ for $N = 2$. So the Higgs boson will escape from being detected. However for total luminosity $L = 4 \cdot 10^5 \text{ pb}^{-1}$ the signal significance is bigger than $S \geq 5.4$ and 3 Higgs like bosons could be discovered. Similar analysis works for the Higgs boson mass range interval $140 \text{ GeV} \leq m_h \leq 400 \text{ GeV}$, where the most promising signature is the use of the Higgs boson decay into 4 leptons (9). As it follows from the LEP1 data [7] it is possible for the Higgs like bosons with masses $m \leq 30 \text{ GeV}$ to escape from the detection for the maximal mixing case provided $N \geq 10$.

As a toy example consider the following mass spectrum of the $\phi'_i(x)$ fields:

$$M^2(i) = M_0^2 + \frac{\delta \cdot i}{N+1}, \quad (15)$$

where $i = 1, 2, \dots, N+1$. In the limit $N \rightarrow \infty$ we shall have the Higgs boson with continuously distributed mass $M_0^2 \leq M_h^2 \leq M_0^2 + \delta$. For $\delta \sim M^2$ Higgs boson has big decay width

and escapes from the detection. Such Higgs boson with continuously distributed mass looks like broad resonance with decay width $\Gamma = \sqrt{M_0^2 + \delta} - M_0$ and a mass $M_0^2 + \frac{\delta}{2}$. The branching ratios for such "modified" Higgs boson coincide with the corresponding branching ratios of the standard Higgs boson unlike to the model of ref. [6] where the big decay width of the Higgs boson is due to Higgs boson decays into invisible modes. Such models with continuously distributed Higgs boson mass have been discussed in ref. [4].

As a quasirealistic example consider NMSSM(NMSSM = MSSM + chiral $SU(2) \otimes U(1)$ singlet superfield) [10]. The superpotential of the NMSSM contains a term $\delta W = \lambda \sigma \bar{H} H$ in the superpotential that relates the $SU(2) \otimes U(1)$ singlet chiral superfield σ with chiral superhiggs fields H and $\bar{H}$. For small coupling constant λ singlet superfield σ effectively decouples from the MSSM superfields and most of the NMSSM predictions(spectrum, branching ratios, cross sections...) coincide with the corresponding MSSM predictions. The general form of the soft breaking terms for complex scalar singlet field $\sigma(x) = \frac{\sigma_1(x) + i\sigma_2(x)}{\sqrt{2}}$ is

$$V_{soft,\sigma} = \frac{m_1^2 \sigma_1^2}{2} + \frac{m_2^2 \sigma_2^2}{2} + m_{12}^2 \sigma_1 \sigma_2 + (k_1 \sigma_1 + k_2 \sigma_2) \bar{H} H + c_1 \sigma_1 + c_2 \sigma_2 + \dots \quad (16)$$

The nondiagonal terms in (16) lead to nonzero mixing between the singlet fields $\sigma_1(x)$, $\sigma_2(x)$ and the lightest Higgs field $h(x)$. It is possible to choose parameters of the model such that the mixing is the maximal one, i.e.

$$h(x) = \frac{1}{\sqrt{3}}(\phi_1(x) + \phi_2(x) + \phi_3(x)), \quad (17)$$

where the fields $\phi_i(x)$ have masses M_i . As it has been discussed before for such case we shall have Higgs boson signal dilution compared to the standard case.

To conclude, in this paper we have studied the influence of scalar singlets on the Higgs boson detection at LHC. We have found that big mixing of singlet fields with Higgs boson field can lead to the nonobservation of the Higgs boson at LHC or to the observation of the additional Higgs like states.

I am indebted to the collaborators of the INR theoretical department for discussions and critical comments. I thank CERN TH Department for the hospitality during my stay

at CERN where this paper has been finished. The research described in this publication was made possible in part by Award No RP1-187 of the U.S. Civilian Research and Development Foundation for the Independent States of the Former Soviet Union(CRDF).

Note added. After this paper has been finished I became aware of paper by Annindya Datta and Amitava Raychandhuty, hep-ph/9708444, where the effects of mixing between singlet and Higgs fields have been studied.

References

- [1] A.Sopczak. Talk given at the international conference
Non-Accelerator searches for new physics, Dubna, 7-11 july 1997.
- [2] B.W.Lee, C.Quigg and C.B.Thacker, Phys.Rev. **D10**(1974)1145.
- [3] J.Ellis, Phenomenology of LEP2 physics, CERN-TH/97-131.
- [4] N.V.Krasnikov, Phys.Lett.**B325**(1994)430.
- [5] R.E.Schrock and M.Suzuki, Phys.Lett.**B100**(1982)250;
L.F.Li, Y.Liu and L.Wolfenstein, Phys.Lett.**B159**(1985)45;
D.Chang and W.Keung, Phys.Lett.**B217**(1989)238;
N.V.Krasnikov, Phys.Lett.**B291**(1992)89;
A.S. Joshipura and J.W.F.Valle, Nucl.Phys.**B397**(1993)105.
- [6] T.Binoth and J.J. van der Bij, Z.Phys.**C75**(1997)17.
- [7] O.Adriani et al., Phys.Lett.**B294**(1992)457;
D.Buskulic et al., Phys.Lett.**B313**(1993)312.
- [8] CMS, Technical Proposal, CERN/LHCC/94-38 LHCCP1, 15 december 1994.
- [9] ATLAS, Technical Proposal. CERN/LHCC/94-43 LHCCP2, 15 december 1994.
- [10] J.Ellis et al., Phys.Rev.**D39**(1989)844;
L.Durand and J.L.Lopez, Phys.Lett.**B217**(1989)463;
S.F.King and P.L.White, Phys.Rev.**D52**(1995)4183;
M.Drees, Int.J. Mod.Phys.**A4**(1989)3635;
U.Ellwanger, M.Rausch de Traubenberg and C.A.Savoy, Phys.Lett**B315**(1993)331;
Z.Phys.**C67**(1995)665; Nucl.Phys.**B492**(1997)21.